


Biotage® PS-Triphenylphosphine

Solid-Supported Phosphine



Key Facts



Stoichiometric



Shelf Life

Capacity
(mmol/g)

BSE/TSE



Scalable

Particle Size
(μm)Thermally &
Mechanically
StableGood
Laboratory
PracticeBulk Density
(g/L)


Specifications

Chemical Name:	Diphenylphosphino-polystyrene
Resin Type:	1% Cross-linked poly(styrene-co-divinylbenzene)
Application:	Chlorination of acids and alcohols, Wittig and Mitsunobu reactions, scavenging of alkyl halides
Typical Chlorination Conditions:	0.5 equivalent of acid or alcohol in CCl_4 , 3 h, reflux
Typical Mitsunobu Reaction Conditions:	1.0 equivalent of alcohol, 1.5 equivalent of phenol, 2.2 equivalents of resin, and 1.6 equivalent of di-tert-butyl azodicarboxylate (DBAD) at room temperature for 16 h
Typical Wittig Reaction Conditions:	2.0 equivalents of ylide resin, 8.0 equivalent of sodium bis(dimethylsilyl)amide/tetrahydrofuran (NaHMDS/THF), resin washed with THF, followed by 1.0 equivalent of carbonyl compound in THF at room temperature for 16 h
Typical Alkyl Halide Scavenging Conditions:	3.0 equivalent of resin, DMF, 10 mL/g resin, 20 °C, 16 h

Biotage® PS-Triphenylphosphine is a diphenylphosphinated polystyrene resin that is a solid-supported equivalent of triphenylphosphine. The capacity of the resin is determined by the quantitation of benzyl bromide uptake in DMF (GC, internal standard method). The resin can readily convert

alcohols or carboxylic acids to the corresponding chlorides or acid chlorides in carbon tetrachloride (Scheme 1).¹⁻³ Reaction conditions are relatively mild and the products are formed in high yield and purity.


PS-Triphenylphosphine can also be used in Mitsunobu reactions to prepare aryl ethers in good-to-excellent yields and in high purities (Scheme 2).⁴ The procedures described provide pure products without the need for laborious silica gel chromatography. Removal of excess phenol is accomplished



Scheme 1. Chlorination using PS-Triphenylphosphine.

with MP-Carbonate, while removal of excess DBAD (di-tert-butyl azodicarboxylate)-based hydrazide by-products may be accomplished by addition of TFA, followed by separation with silica or by liquid-liquid extraction (LLE). Reactions may be carried out at room temperature. Comparative studies suggest that there is no significant advantage to degassing the reaction mixture with nitrogen contrary to previous recommendations.⁶ However, the order of addition of the substrates is critical in minimization of side products.


PS-Triphenylphosphine resin can also be used to synthesize olefins via the Wittig reaction (Scheme 3)^{5,6} or as a scavenger for alkyl halides (Scheme 4).



Scheme 2. Mitsunobu reaction using PS-Triphenylphosphine.

Application Areas:

- » Mitsunobu chemistry - in the synthesis of aryl or alkyl ethers¹
- » Chlorination - for the ready conversion of alcohols and acids to the corresponding halides or acid chlorides²
- » Wittig chemistry - for C=C double bond formation in the synthesis of olefins³
- » Scavenging - of alkyl halides in solution
- » Cyanation - of aryl halides to form aryl cyanides⁴
- » Cyclisation (condensation) to form 1,3,4-oxadiazoles⁵



Representative Procedures

Mitsunobu Reaction (Table 1, Entry 3)


To a reaction vessel containing PS-Triphenylphosphine resin (311 mg, 0.66 mmol) was added a solution of p-methoxyphenol (0.45 mmol in 1 mL anhydrous THF). The suspension was allowed to stand for 5 minutes and then a solution of DBAD (di-tert-butyl azodicarboxylate) (1 mL, 0.48 mmol in anhydrous THF) was added. A further 0.5 mL of THF was added and the solution agitated at room temperature for 30 min. A solution of benzyl alcohol (1 mL, 0.3 mmol in anhydrous THF) was added and the reaction stirred overnight. In order to scavenge any excess phenols, MP-Carbonate resin (274 mg, 0.75 mmol) was then added and the mixture was stirred for a further 2 h. The resin was filtered and washed with THF (2 x 2 mL). To the filtrate was added 5 mL of a TFA/DCM/water (50:48:2) solution, and the mixture was stirred at RT for 2 h. The product was then extracted with MTBE, washed with water, and the MTBE layer concentrated to afford the product in 81% yield and 100% purity.

Chlorination (Table 2, Entry 4)

To a suspension of PS-Triphenylphosphine resin in CCl₄ (1 g, 2.12 mmol in 8 mL) was added a solution of piperonyl alcohol in CCl₄ (152 mg, 1 mmol in 2 mL). The reaction was heated at reflux for 3 h after which the mixture was filtered and the filtrate concentrated to give pure piperonyl chloride in 100% theoretical yield and purity.

Wittig Reaction (Table 3, Entry 2)

1-Iodobutane (0.53 mL, 8.52 mmol) was added to a suspension of PS-Triphenylphosphine (3.0 g, 4.26 mmol) in 30 mL DMF and the reaction was stirred for 48 h at 65 °C. The resulting phosphonium resin was washed with DMF (4 x 40 mL), toluene (4 x 40 mL), DCM (4 x 40 mL), and diethyl ether (4 x 40 mL) and dried in vacuo for 12 h. The dried phosphonium resin (0.2 g, 0.2 mmol) was added to a reaction vessel, followed by the addition of THF (2 mL). To the suspension of phosphonium resin was added a solution of sodium bis(dimethylsilyl)amide (2.0 M NaHMDS in THF, 0.4 mL, 0.8 mmol) at room temperature and the reaction stirred for 1 h. The ylide resin was washed with THF (5 x 4 mL) to remove excess base. To the suspension of ylide resin in anhydrous THF (2 mL) was added a solution of p-methoxybenzaldehyde (0.2 mL, 0.1 mmol) in THF (2 mL) and the mixture was stirred for 16 h. The reaction mixture was diluted with 2 mL hexane and directly applied to a silica SPE cartridge⁸ followed by washing with hexane/ether (2:1, 2 x 4 mL). The solvent was concentrated to provide the olefin in 94% yield (GC purity 91%).



Scheme 3. Wittig reaction using PS-Triphenylphosphine.





































Entry	Alcohol	Phenol	Aryl Ether	% Yield	% Purity
1				80	100
2				73	98
3				81	100
4				87	97
5				92	100
6				84	100
7				81	100
8				76	100
9				91	96
10				86	100 ^a
11				94	100 ^a
12				92	100 ^a

Table 1. Mitsunobu reaction using PS-Triphenylphosphine resin.

^aConversion 100%. The product was comprised of a mixture of hydrolyzed and un-hydrolyzed acetal protecting group. The product may be purified by LLE using aqueous base and MTBE.













Entry	Alcohol	Product	% Isolated Yield	% Purity ^a
1			98	95
2			74 ^b	64 ^c
3			100	100
4			100	100
5			100	100
6			73	95

Table 2. Chlorination of acids and alcohols using PS-Triphenylphosphine.

^aGC analysis: HP-5 phenylmethylsilicone column 100–250 °C, 15 °C/min, 10 min hold.

^bNot isolated.

^cRate of reaction slower for hindered aliphatic secondary alcohols, 28% conversion after 3 h, 64% conversion after 16 h reflux. Determined by ¹H NMR.










Entry	Phosphonium Resin	Carbonyl Compound	Olefin	% Isolated Yield (cis:trans) ^a	% GC Purity ^b
1				81 (5:1)	95
2				94 (2:3)	91
3				88 (2:1)	94


Table 3. Wittig reaction using PS-Triphenylphosphine.

^aRatio determined by ¹H NMR analysis.

^bGC analysis: HP-5 phenylmethylsilicone column 100–250 °C, 15 °C/min, 10 min hold.

Scavenging of Alkyl Halide (Table 4, Entry 2)

PS-Triphenylphosphine (3.0 equivalents) was added to a solution of benzyl bromide (1.0 equivalent) in DMF (10 mL/g resin added) and the reaction stirred at room temperature for 5–16 h. Results by GC analysis indicate >80% scavenging after 6 h and 100% scavenging after 16 h.



Scheme 4. Scavenging of alkyl halides using PS-Triphenylphosphine

Entry	Material Scavenged	Solvent	Temp °C	% Scavenged ^a	Time (h)
1	Ethyl bromoacetate	DMF	20	100	6 h
2	Benzyl bromide	DMF	20	100	16 h
3	Cinnamyl chloride	THF:DMF	50	100	10 h
4	Cinnamyl bromide	DMF	20	100	16 h

Table 4. Scavenging of alkyl halides with PS-Triphenylphosphine resin (3.0 equiv.)
^aGC analysis: HP-5 phenylmethylsilicone column 100–250 °C, 15 °C/min, 10 min hold.

Ordering Information

Part Number	Quantity
800510	3 g
800378	10 g
800379	25 g
800380	100 g
800381	1000 g

References

- Relles, H. M.; Schluenz, R. W. *J. Am. Chem. Soc.* 1974, 96, 6469.
- Regen, S. L.; Lee, D. P. *J. Org. Chem.* 1975, 40, 1669.
- Landi, J. J. Jr.; Brinkman, H. R. *Synthesis* 1992, 1093.
- Tunoori, A. R.; Dutta, D.; Georg, G. I. *Tetrahedron Lett.* 1998, 39, 8951.
- Vasudevan, A.; Park, D. C.; Wodka, D.; Gentles, R. G. *J. Comb. Chem.* 2002, 4, 442.
- Bernard, M.; Ford, W. T. *J. Org. Chem.* 1983, 48, 326.
- Bolli, M. H.; Ley, S. V. *J. Chem. Soc., Perkin Trans. 1.* 1998, 15, 2243.
- Part Number 440-0100-C. Compatible Solvents: DMF (3.5 mL/g), THF (4.1 mL/g), DCM (5 mL/g), benzene (3.1 mL/g).



FM 31206 EMS 640981

Biotage holds certification for both ISO9001 Quality Management and ISO14001 Environmental Management.

EUROPE

Main Office: +46 18 565900
 Toll Free: +800 18 565710
 Fax: +46 18 591922
 Order Tel: +46 18 565710
 Order Fax: +46 18 565705
 order@biotage.com
 Support Tel: +46 18 56 59 11
 Support Fax: + 46 18 56 57 11
 eu-1-pointsupport@biotage.com

NORTH & LATIN AMERICA

Main Office: +1 704 654 4900
 Toll Free: +1 800 446 4752
 Fax: +1 704 654 4917
 Order Tel: +1 704 654 4900
 Order Fax: +1 434 296 8217
 ordermailbox@biotage.com
 Support Tel: +1 800 446 4752
 Outside US: +1 704 654 4900
 us-1-pointsupport@biotage.com

JAPAN

Tel: +81 3 5627 3123
 Fax: +81 3 5627 3121
 jp_order@biotage.com
 jp-1-pointsupport@biotage.com

CHINA

Tel: +86 21 2898 6655
 Fax: +86 21 2898 6153
 cn_order@biotage.com
 cn-1-pointsupport@biotage.com

To locate a distributor, please visit our website www.biotage.com

Part Number: PPS389.V.1

© 2016 Biotage. All rights reserved. No material may be reproduced or published without the written permission of Biotage. Information in this document is subject to change without notice and does not represent any commitment from Biotage. E&OE. A list of all trademarks owned by Biotage AB is available at www.biotage.com/legal. Other product and company names mentioned herein may be trademarks or registered trademarks and/or service marks of their respective owners, and are used only for explanation and to the owners' benefit, without intent to infringe.